




CHEMISTRY MARKING SCHEME


SET -56/3


Compt. July, 2015



Qu es.	Value points	Marks
1	Formation of stable complex by polydentate ligand.	1
2	Propanal	1
3	p-Nitroaniline < Aniline < p-Toluidine	1
4	Frenkel defect	1
5	Emulsions are liquid – liquid colloidal systems. For example – milk, cream (or any other one correct example)	½ + ½
6	Potassium permanganate is prepared by fusion of MnO ₂ with an alkali metal hydroxide and an oxidising agent like KNO ₃ . This produces the dark green K ₂ MnO ₄ which disproportionates in a neutral or acidic solution to give permanganate. 2MnO₂ + 4KOH + O₂ → 2K₂MnO₄ + 2H₂O 3MnO₄²⁻ + 4H⁺ → 2MnO₄⁻ + MnO₂ + 2H₂O Oxalate ion or oxalic acid is oxidised at 333 K: 5C₂O₄²⁻ + 2MnO₄⁻ + 16H⁺ → 2Mn²⁺ + 8H₂O + 10CO₂ OR	1 1
6	i) Iodine is liberated from potassium iodide : 10I⁻ + 2MnO₄⁻ + 16H⁺ → 2Mn²⁺ + 8H₂O + 5I₂ ii) Hydrogen sulphide is oxidised, sulphur being precipitated: H₂S → 2H⁺ + S²⁻ 5S²⁻ + 2MnO₄⁻ + 16H⁺ → 2Mn²⁺ + 8H₂O + 5S	1 1
7	<p>The mechanism shows the following steps:</p> $\text{H}-\underset{\text{H}}{\overset{\text{H}}{\text{C}}}-\underset{\text{H}}{\overset{\text{H}}{\text{C}}}-\ddot{\text{O}}-\text{H} + \text{H}^+ \xrightleftharpoons{\text{Fast}} \text{H}-\underset{\text{H}}{\overset{\text{H}}{\text{C}}}-\underset{\text{H}}{\overset{\text{H}}{\text{C}}}-\overset{\text{H}}{\text{O}^+}-\text{H}$ $\text{H}-\underset{\text{H}}{\overset{\text{H}}{\text{C}}}-\underset{\text{H}}{\overset{\text{H}}{\text{C}}}-\overset{\text{H}}{\text{O}^+}-\text{H} \xrightleftharpoons{\text{Slow}} \text{H}-\underset{\text{H}}{\overset{\text{H}}{\text{C}}}-\overset{\text{H}}{\text{C}^+} + \text{H}_2\text{O}$ $\text{H}-\underset{\text{H}}{\overset{\text{H}}{\text{C}}}-\overset{\text{H}}{\text{C}^+} \rightleftharpoons \text{H}-\underset{\text{H}}{\text{C}}=\underset{\text{H}}{\text{C}}-\text{H} + \text{H}^+$ <p align="center">Ethene</p>	½ ½ 1

8	<p>i) Mole fraction of a component =</p> $\frac{\text{Number of moles of the component}}{\text{Total number of moles of all the components}}$ <p>ii) Molality (<i>m</i>) is defined as the number of moles of the solute per kilogram (kg) of the solvent.</p> <p style="text-align: center;">Or</p> $\text{Molality (m)} = \frac{\text{Moles of solute}}{\text{Mass of solvent in kg}}$	1 1
9	<p>Zero order : mol L⁻¹s⁻¹</p> <p>Second order : L mol⁻¹s⁻¹</p>	1 1
10	<p>i) Due to high bond dissociation enthalpy of N ≡ N</p> <p>ii) Due to low bond dissociation enthalpy of F₂ than Cl₂ and strong bond formation between N and F</p>	1 1
11	<p>Disproportionation : The reaction in which an element undergoes self-oxidation and self-reduction simultaneously. For example –</p> $2\text{Cu}^+(\text{aq}) \longrightarrow \text{Cu}^{2+}(\text{aq}) + \text{Cu}(\text{s})$ <p>(Or any other correct equation)</p>	1 ½ 1 ½
12	<p>i) Hexaamminecobalt(III) chloride</p> <p>ii) Tetrachlorido nickelate(II)</p> <p>iii) Potassium hexacyanoferrate(III)</p>	1 1 1
13	<p>i) 2-bromobutane</p> <p>ii) 1, 3-dibromobenzene</p> <p>iii) 3-choloropropene</p>	1 1 1
14	<p>i) </p> <p>ii) $\text{CH}_3\text{CH}_2\text{MgCl} \xrightarrow[\text{H}_2\text{O}]{\text{HCHO}} \text{CH}_3\text{-CH}_2\text{-CH}_2\text{-OH}$</p> <p>$\text{CH}_3\text{CH}=\text{CH}_2 + \text{H}_2\text{O} \xrightleftharpoons{\text{H}^+} \text{CH}_3\text{-}\underset{\text{OH}}{\text{CH}}\text{-CH}_3$</p>	1 1 1
15	<p>i) $\text{CH}_3\text{-CH}_2\text{OH} \xrightarrow{\text{PCl}_5} \text{CH}_3\text{CH}_2\text{Cl}$</p>	1

	<p>ii)</p>  <p>iii)</p> $\text{CH}_3\text{Cl} + \text{CH}_3\text{CH}_2\text{-ONa} \longrightarrow \text{CH}_3\text{CH}_2\text{-O-CH}_3$	<p>1</p> <p>1</p>
16	<p>i) Peptide linkage – in proteins, α-amino acids are connected to each other by peptide bond or peptide linkage (-CONH- bond).</p> <p>ii) Primary structure - each polypeptide in a protein molecule having amino acids which are linked with each other in a specific sequence.</p> <p>iii) Denaturation - When a protein is subjected to physical change like change in temperature or chemical change like change in pH, protein loses its biological activity.</p>	<p>1</p> <p>1</p> <p>1</p>
17	<p>Copolymerisation is a polymerisation reaction in which a mixture of more than one monomeric species is allowed to polymerise and form a copolymer.</p>  <p>1,3-Butadiene + Styrene \longrightarrow Butadiene - styrene copolymer</p> <p>1,3-Butadiene + Acrylonitrile $\xrightarrow{\text{Copolymerisation}}$ Buna-N (or any other correct example)</p>	<p>1</p> <p>1</p> <p>1</p>
18	$r = \frac{\sqrt{2}a}{4}$ $r = \frac{1.414 \times 4.077 \times 10^{-8} \text{ cm}}{4}$ $r = 1.44 \times 10^{-8} \text{ cm}$	<p>1</p> <p>1</p> <p>1</p>
19	$\pi_{\text{cane sugar}} = \pi_X$ <p>Therefore, $c_{\text{cane sugar}} = c_X$ (where c is molar concentration)</p> $\frac{W_{\text{cane sugar}}}{M_{\text{cane sugar}}} = \frac{W_X}{M_X}$ $\frac{5 \text{ g}}{342 \text{ g mol}^{-1}} = \frac{0.877}{M_X}$ $M_X = \frac{0.877 \times 342}{5} \text{ gmol}^{-1}$ $M_X = 59.9 \text{ or } 60 \text{ gmol}^{-1}$	<p>1</p> <p>1</p> <p>1</p>
20	$k = \frac{2.303}{t} \log \frac{[R]_0}{[R]}$	<p>1</p>

	$60 \text{ s}^{-1} = \frac{2.303}{t} \log \frac{[R]_0}{\frac{[R]_0}{10}}$ $t = \frac{2.303}{60 \text{ s}^{-1}} \log 10$ $t = \frac{2.303}{60} \text{ s}$ $t = 0.0384 \text{ s}$	1 1				
21	<p>i) It is a process of removing the dissolved substance from a colloidal solution by means of diffusion through a semi - permeable membrane.</p> <p>ii) The movement of colloidal particles under an applied electric potential towards oppositely charged electrode is called electrophoresis.</p> <p>iii) Colloidal particles scatter light in all directions in space. This scattering of light illuminates the path of beam in the colloidal dispersion.</p>	1 1 1				
22	<p>i) It lowers the melting point of alumina / acts as a solvent.</p> <p>ii)</p> <table border="1" style="margin-left: auto; margin-right: auto;"> <thead> <tr> <th>Roasting</th> <th>Calcination</th> </tr> </thead> <tbody> <tr> <td>Ore is heated in a regular supply of air</td> <td>Heating in a limited supply or absence of air.</td> </tr> </tbody> </table> <p>(Or with equation)</p> <p>iii) It is a process of separation of different components of a mixture which are differently adsorbed on a suitable adsorbent.</p> <p style="text-align: center;">OR</p>	Roasting	Calcination	Ore is heated in a regular supply of air	Heating in a limited supply or absence of air.	1 1 1
Roasting	Calcination					
Ore is heated in a regular supply of air	Heating in a limited supply or absence of air.					
22	<p>$3\text{Fe}_2\text{O}_3 + \text{CO} \rightarrow 2\text{Fe}_3\text{O}_4 + \text{CO}_2$ (Iron ore)</p> <p>$\text{Fe}_3\text{O}_4 + \text{CO} \rightarrow 3\text{FeO} + \text{CO}_2$</p> <p>$\text{CaCO}_3 \rightarrow \text{CaO} + \text{CO}_2$ (Limestone)</p> <p>$\text{CaO} + \text{SiO}_2 \rightarrow \text{CaSiO}_3$ (Slag)</p> <p>$\text{FeO} + \text{CO} \rightarrow \text{Fe} + \text{CO}_2$</p> <p>$\text{C} + \text{CO}_2 \rightarrow 2\text{CO}$ Coke</p> <p>$\text{C} + \text{O}_2 \rightarrow \text{CO}_2$</p> <p>$\text{FeO} + \text{C} \rightarrow \text{Fe} + \text{CO}$</p> <p style="text-align: right;">(any 6 correct equations)</p>	$6 \times \frac{1}{2}$ $= 3$				
23	<p>i) Aspartame, Saccharin (any one)</p> <p>ii) No</p> <p>iii) Social concern, empathy, concern, social awareness (any 2)</p>	1 1 2				
24	<p>a) i)</p>  <p>ii)</p> <p>$(\text{CH}_3)_2\text{C}=\text{CHCOCH}_3$</p> <p>b) i) Add NaHCO_3, benzoic acid will give brisk effervescence of CO_2 whereas ethylbenzoate</p>	1 1				

	<p>will not.</p> <p>ii) Add NaOH and I₂, acetophenone forms yellow ppt of iodoform on heating whereas benzaldehyde will not.</p> <p>iii) Add neutral FeCl₃, phenol gives violet colouration whereas benzoic acid does not.</p> <p style="text-align: right;">(or any other correct test)</p> <p style="text-align: center;">OR</p>	<p>1</p> <p>1</p> <p>1</p>
24	<p>a) i)</p> $\begin{array}{c} \text{CH}_3 \\ \diagdown \\ \text{C}=\text{N}-\text{OH} \\ \diagup \\ \text{CH}_3 \end{array}$ <p>ii)</p> $\begin{array}{c} \text{CH}_3 \\ \diagdown \\ \text{C}=\text{N}-\text{NH}-\overset{\text{O}}{\parallel}{\text{C}}-\text{NH}_2 \\ \diagup \\ \text{H} \end{array}$ <p>b) i)</p> $\text{CH}_3\text{CHO} \xrightarrow[\text{conc HCl}]{\text{Zn-Hg}} \text{CH}_3-\text{CH}_3$ <p>ii)</p> $2 \text{CH}_3-\text{CHO} \xrightleftharpoons{\text{dil. NaOH}} \text{CH}_3-\underset{\text{OH}}{\text{CH}}-\text{CH}_2-\text{CHO}$ <p>iii)</p> $\text{CH}_3\text{CHO} \xrightarrow{\text{LiAlH}_4} \text{CH}_3\text{CH}_2\text{OH}$	<p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p>
25	<p>a) Due to relatively stable half – filled p-orbitals of group 15 elements</p> <p>b) i) $\text{CaF}_2 + \text{H}_2\text{SO}_4 \rightarrow \text{CaSO}_4 + 2\text{HF}$</p> <p>ii) $\text{SO}_2(\text{g}) + \text{Cl}_2(\text{g}) \rightarrow \text{SO}_2\text{Cl}_2(\text{l})$</p> <p>iii) $2\text{NH}_4\text{Cl} + \text{Ca}(\text{OH})_2 \rightarrow 2\text{NH}_3 + 2\text{H}_2\text{O} + \text{CaCl}_2$</p> <p style="text-align: center;">OR</p>	<p>2</p> <p>1</p> <p>1</p> <p>1</p>
25	<p>a) i)</p> 	<p>1</p>

	<p>ii)</p>  <p>b) i) Due to small size of nitrogen, the lone pair of electron on nitrogen is localized/ easily available for donation. ii) Because they need only one electron to attain stable/noble gas configuration.</p>  <p>iii)</p>	<p>1</p> <p>1</p> <p>1</p> <p>1</p>
26	$E^0_{\text{cell}} = E^0_{\text{Sn}^{2+}/\text{Sn}} - E^0_{\text{Zn}^{2+}/\text{Zn}}$ $= -0.14\text{V} - (-0.76\text{V})$ $= 0.62\text{V}$ $\Delta_r G^0 = -n F E^0_{\text{cell}}$ $= -2 \times 96500 \text{ C mol}^{-1} \times 0.62 \text{ V}$ $= -119660 \text{ J mol}^{-1}$ $E_{\text{cell}} = E^0_{\text{cell}} - \frac{0.059}{n} \log \frac{[\text{Zn}^{2+}]}{[\text{Sn}^{2+}]}$ $E_{\text{cell}} = 0.62 - \frac{0.059}{2} \log \frac{[\text{Zn}^{2+}]}{[\text{Sn}^{2+}]}$ <p style="text-align: center;">OR</p>	<p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p>
26	<p>a) The conductivity of a solution at any given concentration is the conductance of one unit volume of solution kept between two platinum electrodes with unit area of cross section and at a distance of unit length. Molar conductivity of a solution at a given concentration is the conductance of the volume V of solution containing one mole of electrolyte kept between two electrodes with area of cross section A and distance of unit length. Molar conductivity increases with decrease in concentration.</p> <p>b) $E^0_{\text{cell}} = E^0_{\text{C}} - E^0_{\text{A}}$ $= 0.80\text{V} - 0.77\text{V}$ $= 0.03\text{V}$ $\Delta_r G^0 = -n F E^0_{\text{cell}}$ $= -1 \times 96500 \text{ C mol}^{-1} \times 0.03 \text{ V}$ $= -2895 \text{ J mol}^{-1}$ $\text{Log } K_c = \frac{n E^0_{\text{cell}}}{0.059}$</p>	<p>1/2</p> <p>1/2</p> <p>1</p> <p>1/2</p> <p>1/2</p> <p>1</p> <p>1/2</p>

$\text{Log } K_c = \frac{1 \times 0.03}{0.059}$ $\text{Log } K_c = 0.508$	$\frac{1}{2}$
---	---------------

Dr. Sangeeta Bhatia

Sh. S.K. Munjal

Sh. D.A. Mishra

Ms. Garima Bhutani