














CHEMISTRY MARKING SCHEME
2015
56/1/RU




. NO.	Value points	MARKS
Q.1	H_2SO_3 H_2SO_4 $\text{H}_2\text{S}_2\text{O}_8$, H_2SO_5 (any two formulae)	$\frac{1}{2} + \frac{1}{2}$
Q.2	1-ethoxy-2-methylpropane	1
Q.3	Due to coagulation of colloidal clay particles	1
Q.4	$\text{CH}_3\text{-CH(Br)-CH}_3$	1
Q.5	X_4Y_3	1
Q.6	Similarity : Both show contraction in size /Both show irregularity in their electronic configuration/Both are stable in +3oxidation state (any one) Difference :Actinoids are mainly radioactive but lanthanoids are not/ Actinoids show wide range of oxidation states but lanthanoids do not /Actinoid contraction is greater than lanthanoid contraction. (any other one similarity and one difference)	1 1
Q.7	(i) Pentaamminechloridocobalt(III) ion (ii) $\text{K}_2[\text{NiCl}_4]$	1 1
Q.8	(i) PCC / Cu at 573 K (ii) NH_3 , Δ (heat) OR 8. (i) $\text{C}_6\text{H}_5\text{COCH}_3 < \text{CH}_3\text{COCH}_3 < \text{CH}_3\text{CHO}$ (ii) $\text{CH}_3\text{COOH} < \text{Cl-CH}_2\text{-COOH} < \text{F-CH}_2\text{-COOH}$	1 1 1 1
Q.9	(i) Negative deviation ,temperature will increase. (ii) Blood cell will swell due to osmosis , water enters into the cell.	$\frac{1}{2} + \frac{1}{2}$ $\frac{1}{2} + \frac{1}{2}$
Q.10	$\text{Cu}^{2+} + 2\text{e} \rightarrow \text{Cu}$ 63.5 g Cu is deposited = 2×96500 C 1.27 g Cu is deposited = $2 \times 96500 \times 1.27 / 63.5$ C = ixt (Q = ixt) $t = 2 \times 96500 \times 1.27 / 63.5 \times 2 = 1930\text{s}$ Or	1 1

	<p>by Faraday First law</p> $m = z x i x t$ <p>$z = \text{atomic mass/valency} \times F$</p> $1.27 = 63.5 \times 2 x t / 2 \times 96500$ $t = 1930 \text{ s}$	<p>½</p> <p>½</p> <p>1</p>
Q.11	$\frac{p^0 - p}{p^0} = \frac{w_s \times M_{\text{solvent}}}{M_s \times W_{\text{solvent}}}, \quad s = \text{solute}$ $(32 - 31.84) / 32 = 10 \times 18 / M_s \times 200$ $M_s = 180 \text{ g/mol}$	<p>1</p> <p>1</p> <p>1</p>
Q.12	<p>(i) Zone refining</p> <p>(ii) SiO_2 act as flux to remove the impurity of Iron oxide</p> <p>(iii) Depressants prevent one type of sulphide ore forming the froth with air bubbles.</p>	<p>1</p> <p>1</p> <p>1</p>
Q.13	<p>(i) Starch.</p> <p>(ii) α- Helix polypeptide chains are stabilized by intramolecular H-bonding whereas β- pleated sheet is stabilized by intermolecular H-bonding. (or any other difference)</p> <p>(iii) Pernicious anaemia</p>	<p>1</p> <p>1</p> <p>1</p>
Q.14	<p>(i) Hydration isomerism</p> <p>(ii) Electronic configuration $1s^2 2s^2 2p^6 3s^2 3p^4$ / or by diagram</p> <p>(iii) Hybridization is $sp^3 d^2$ and shape is octahedral.</p>	<p>1</p> <p>1</p> <p>½ + ½</p>
Q.15	<p>(i)</p> <div style="text-align: center;"> <p style="text-align: center;">Benzene diazonium halide</p> </div> <div style="text-align: center;"> <p style="text-align: right;">(where X=Br)</p> </div> <p>(ii)</p>	<p>1</p>

15	 <p>iii) $\text{CH}_3\text{CH}_2\text{Cl} \xrightarrow[\text{dry ether}]{\text{Na}} \text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_3$</p> <p style="text-align: center;">OR</p> <p>(i) </p> <p>(ii) $\text{CH}_3\text{CH}_2\text{Cl} + \text{AgNO}_2 \rightarrow \text{CH}_3\text{CH}_2\text{NO}_2 + \text{AgCl}$</p> <p>(iii) $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}(\text{Br})\text{CH}_3 + \text{KOH (alc.)} \rightarrow \text{CH}_3\text{CH}_2\text{CH}=\text{CHCH}_3$</p>	1 1 1 1 1
Q.16	<p>(i) Stoichiometric defect</p> <p>(ii) Schottky defect e.g. NaCl (or any other example)</p> <p>(iii) Density of crystal decreases</p>	1 $\frac{1}{2} + \frac{1}{2}$ 1
Q.17	$\Lambda_m = \frac{1000 \times k}{M} \text{ Scm}^2\text{mol}^{-1}$ $\Lambda_m = \frac{1000 \times 5.25 \times 10^{-5}}{2.5 \times 10^{-4}} \text{ Scm}^2\text{mol}^{-1}$ $= 210 \text{ Scm}^2\text{mol}^{-1}$ $\Lambda_m^0 \text{ HCOOH} = \lambda^0 \text{ HCOO}^- + \lambda^0 \text{ H}^+$ $(50.5 + 349.5) \text{ S cm}^2\text{mol}^{-1} = 400 \text{ S cm}^2\text{mol}^{-1}$ $\alpha = \Lambda_m / \Lambda_m^0$	$\frac{1}{2}$ 1 $\frac{1}{2}$

	$\alpha = 210 / 400 = 0.525$	1
Q.18	<p>Physisorption : adsorbate is held by weak van der Waals' force non-specific It forms multimolecular layer</p> <p>Chemisorption : adsorbate molecules are held by strong forces like a chemical bond It is specific It forms unimolecular layer (or any correct three points)</p>	1,1,1
Q.19	<p>(i) Phenoxide ion is stabilized by resonance as compared to CH_3O^- / In phenol, oxygen acquires +ve charge due to resonance and releases H^+ ion easily whereas there is no resonance in methanol.</p> <p>(ii) Due to lone pair- lone pair repulsion on oxygen.</p> <p>(iii) $(\text{CH}_3)_3\text{C}^+$ is 3^0 carbo-cation which is more stable than CH_3^+ for $\text{S}_{\text{N}}1$ reaction.</p>	1 1 1
Q.20	<p>i) $(\text{CH}_3)_2\text{C}=\text{N}-\text{NH}_2$ ii)  / benzoic acid iii)  / m-bromobenzoic acid</p>	1+1+1
Q.21	<p>(a)</p> <p>(i) Because Cu^+ undergoes disproportionation as $2\text{Cu}^+ \rightarrow \text{Cu} + \text{Cu}^{2+}$</p> <p>(ii) Because of small size of metal, high ionic charge and availability of vacant d-orbital.</p> <p>(b) $\text{Cr}_2\text{O}_7^{2-} + 8\text{H}^+ + 3\text{NO}_2^- \rightarrow 2\text{Cr}^{3+} + 3\text{NO}_3^- + 4\text{H}_2\text{O}$ (Balanced equation only)</p>	1 1 1
Q.22	<p>(i) ethylene glycol $\text{HO}-\text{CH}_2-\text{CH}_2-\text{OH}$</p> <p>Terephthalic acid </p> <p>(ii) 1,3-butadiene $\text{CH}_2=\text{CH}-\text{CH}=\text{CH}_2$</p> <p></p> <p>Styrene</p> <p>(iii) Chloroprene $\text{CH}_2=\text{C}(\text{Cl})-\text{CH}=\text{CH}_2$</p> <p>(Note: Half mark for name/s and half mark for structure/s in each case)</p>	$\frac{1}{2} + \frac{1}{2}$ $\frac{1}{2} + \frac{1}{2}$ $\frac{1}{2}, \frac{1}{2}$
Q.23	(i) Social awareness ,Health conscious, Caring , empathy, concern .(or any other two values)	$\frac{1}{2}, \frac{1}{2}$

	(ii) Cartoon display / street display/poster making (or any other correct answer)	1
	(iii) Wrong choice and over dose may be harmful.	1
	(iv) Saccharin , Aspartame (or any other example)	$\frac{1}{2} + \frac{1}{2}$
Q.24	(a) (i) Due to decrease in bond dissociation enthalpy from HF to HI , there is an increase in acidic character observed. (ii)Oxygen exists as diatomic O ₂ molecule while sulphur as polyatomic S ₈ (iii)Due to non- availability of d orbitals (b)	1 1 1
	<div style="display: flex; justify-content: space-around; align-items: center;"> <div style="text-align: center;">  <p>i)</p> </div> <div style="text-align: center;">  <p>ii)</p> </div> </div> <p style="text-align: center;">OR</p>	1+1
24	(i) White Phosphorus, because it is less stable due to angular strain (ii)Nitrogen oxides emitted by supersonic jet planes are responsible for depletion of ozone layer. Or $\text{NO} + \text{O}_3 \rightarrow \text{NO}_2 + \text{O}_2$ (iii)due to small size of F, large inter electronic repulsion / electron- electron repulsion among the lone pairs of fluorine (iv)Helium (v) $\text{XeF}_2 + \text{PF}_5 \rightarrow [\text{XeF}]^+ [\text{PF}_6]^-$	$\frac{1}{2}, \frac{1}{2}$ 1 1 1 1 1
Q.25	<div style="display: flex; justify-content: space-between; align-items: flex-start;"> <div style="text-align: center;"> <p>A =</p>  </div> <div style="text-align: center;"> <p>B =</p>  </div> <div style="text-align: center;"> <p>C =</p>  </div> </div> <div style="display: flex; justify-content: space-between; align-items: flex-start; margin-top: 20px;"> <div style="text-align: center;"> <p>D =</p>  </div> <div style="text-align: center;"> <p>E =</p>  </div> </div>	1 each

25	<p style="text-align: center;">OR</p> <p>i) </p> <p>ii) </p> <p>iii) </p> <p>(b) $C_2H_5NH_2 < (C_2H_5)_3N < (C_2H_5)_2NH$</p> <p>(c) Add $CHCl_3$ and alc KOH, $C_6H_5-NH_2$ gives foul smell of isocyanide whereas $C_6H_5-NH-CH_3$ does not (or any other correct test)</p>	1,1,1 1 1
Q.26	<p>(a) $[A]_0 = 0.10 \text{ mol/L}$ $[A] = 0.05 \text{ mol/L}$ at time $t = 10s$</p> $k = \frac{2.303}{t} \log \frac{[A_0]}{[A]}$ $k = \frac{2.303}{10 \text{ s}} \log \frac{0.10}{0.05}$ $k = 0.0693s^{-1}$ <p>$t = 20s$</p> $k = \frac{2.303}{t} \log \frac{[A_0]}{[A]}$ $k = \frac{2.303}{20 \text{ s}} \log \frac{0.10}{0.025}$ $k = 0.0693s^{-1}$ <p>As the rate constant is same so it follows pseudo first order reaction.</p> <p>(b) Average rate of reaction = $-\Delta[R]/\Delta t$</p> $= - [0.025 - 0.05 / 20 - 10]$ $= 0.0025 \text{ mol L}^{-1}s^{-1}$	½ 1 1 ½ ½ 1

OR

26

(a)

(i) Rate of reaction becomes 4 times

(ii) Over all order of reaction = 2

1

1

(b)
$$t_{1/2} = \frac{0.693}{k}$$

$$30\text{min} = \frac{0.693}{k}$$

$$k = 0.0231\text{min}^{-1}$$

$$k = \frac{2.303}{t} \log \frac{[A_0]}{[A]}$$

$$t = \frac{2.303 \log \frac{100}{10}}{0.0231}$$

$$t = \frac{2.303 \text{min}}{0.0231}$$

$$t = 99.7\text{min}$$

1

½

½

1