


Strictly Confidential- (For Internal and Restricted Use Only) Secondary School Examination
SUMMATIVE ASSESSMENT - II
March 2015


Marking Scheme – Science (Outside Delhi) 31/3

1. The Marking Scheme provides general guidelines to reduce subjectivity in the marking. It carries only suggested value points for the answer. These are only guidelines and do not constitute the complete answer. Any other individual response with suitable justification should also be accepted even if there is no reference to the text.
2. Evaluation is to be done as per instructions provided in the Marking Scheme. It should not be done according to one's own interpretation or any other consideration. Marking Scheme should be strictly adhered to and religiously followed.
3. If a question has parts, please award marks in the right hand side for each part. Marks awarded for different parts of the question should then be totalled up and written in the left hand margin.
4. If a question does not have any parts, marks be awarded in the left hand side margin.
5. If a candidate has attempted an extra question, marks obtained in the question attempted first should be retained and the other answer should be scored out.
6. Wherever only two/three of a 'given' number of examples/factors/points are expected only the first two/three or expected number should be read. The rest are irrelevant and should not be examined.
7. There should be no effort at 'moderation' of the marks by the evaluating teachers. The actual total marks obtained by the candidate may be of no concern of the evaluators.
8. All the Head Examiners / Examiners are instructed that while evaluating the answer scripts, if the answer is found to be totally incorrect, the (X) should be marked on the incorrect answer and awarded '0' marks.
9. $\frac{1}{2}$ mark may be deducted if a candidate either does not write units or writes wrong units in the final answer of a numerical problem.
10. A full scale of mark 0 to 100 has to be used. Please do not hesitate to award full marks if the answer deserves it.
11. As per orders of the Hon'ble Supreme Court the candidates would now be permitted to obtain photocopy of the Answer Book on request on payment of the prescribed fee. All Examiners/Head Examiners are once again reminded that they must ensure that evaluation is carried out strictly as per value points given in the marking scheme.

MARKING SCHEME
CLASS X – OUTSIDE DELHI

Code No. 31/3

Expected Answer/ Value point SECTION – A		Marks	Total
Q1.	Thirteen	1	1
Q2.	Planaria / hydra / earthworm (any two)	½, ½	1
Q3.	Herbivores	1	1
Q4.		<p style="text-align: right;">Diagram Marking of angles</p>	<p style="text-align: center;">1 ½ × 2 2</p>
Q5.	<p>Management of natural resources – To sustain the resources for future generation and current basic needs of the population.</p> <p>Reuse is a better practice because it can be done at household level with no expense of energy. Reuse does not cause pollution. Eco friendly (any one)</p>	<p style="text-align: center;">1 1 2</p>	<p style="text-align: center;">1 2</p>
Q6.	<p>Biodiversity is the range and number of species of living organisms in a given area.</p> <p>Two advantages- Maintains ecological balance. Helps in maintaining rainfall. Prevent soil erosion. (any two)</p>	<p style="text-align: center;">1 ½, ½</p>	<p style="text-align: center;">2</p>
Q7.	<p>Alkene, C_nH_{2n} Or Alkyne, C_nH_{2n-2}</p> <p>Presence of Ni / Pd / Pt as catalyst</p>	<p style="text-align: center;">½, ½ ½</p>	




		$\frac{1}{2}$	
		$\frac{1}{2}, \frac{1}{2}$	
	Or any other hydrogenation reaction.		3
Q8.	<ul style="list-style-type: none"> • Test 1 (Litmus Test) Take two strips of blue litmus paper. Place a drop each of the alcohol and carboxylic acid on these strips separately. The blue litmus paper turns red in the case of carboxylic acid and remains unaffected in the case of alcohol. • Test 2 (Sodium hydrogen carbonate test / sodium carbonate test) A pinch of sodium hydrogen carbonate or sodium carbonate is added, to both separately. If brisk effervescence with the evolution of a colorless gas is observed, it indicates the presence of carboxylic acid. If no change is observed then it confirms the presence of the alcohol. • Test 3 – Ester test or any other suitable test (any two) 	$\frac{1}{2}$	
		1	
		$\frac{1}{2}$	
		1	3
Q9.	a) D, 2, 8, 8, 1 b) A and E belong to the same group as both have the same no. of valence electrons / same no. of electrons (2) in the outermost shell. c) A and B / D and E. A has a bigger atomic radius than B. or D has a bigger atomic radius than E.	$\frac{1}{2}, \frac{1}{2}$	
		$\frac{1}{2}, \frac{1}{2}$	
		$\frac{1}{2}$	
		$\frac{1}{2}$	3
Q10.	Electronic configuration of element with atomic no. 16 is 2,8,6. Since it has 3 shells, the period no. will be 3. Since the no. of valence electrons is 6, the group no. will be $10 + 6 = 16$. Valency of the element will be $8 - \text{valence electrons}$ ie $8 - 6 = 2$.	1	
		$\frac{1}{2}$	
		$\frac{1}{2}$	
		1	3
Q11.	A – Stigma	1	
	B – Pollen tube	1	
	C – Ovule / germ cell / egg cell	1	3
Q12.	Four categories of contraceptive methods – 1. Barrier method (Condoms) 2. Surgical method or its name 3. Withdrawal method 4. Calendar method 5. Hormonal method 6. IUCD/ Copper T/ Loop (any four)		
	Two advantages –		
	1. Helps in maintaining health of women.	$\frac{1}{2} \times 4$	

	2. Helps in preventing STDs especially AIDS 3. Helps in birth control	(any two)	$\frac{1}{2}, \frac{1}{2}$	3
Q13.	Chromosomes – thread like structures made up of DNA found in the nucleus. The original number of chromosomes becomes half during gamete formation. Hence, when the gametes combine, the original number of chromosomes gets restored in the progeny. (or same thing explained in the form of a flow chart).		1 1,1	 3
Q14.	<ul style="list-style-type: none"> • Speciation – evolution of a new species from pre-existing species • Occurring due to accumulation of variations • By processes like genetic drift / geographical barriers like mountains, rivers etc., leading to incapability to reproduce amongst themselves in the population. <p>Natural selection –</p> <ul style="list-style-type: none"> • Change in frequency of some genes in a population • Which give survival advantage to a species from elimination. • Ex. – in a population of beetles, a new variation (green colour) get survival benefit / advantage to green beetles whereas other (red) perishes. 		$\frac{1}{2} \times 3$ $\frac{1}{2} \times 3$	 3
Q15.	Homologous organs – study of homologous organs suggests that the organs having same structure but performing different functions have evolved from a common ancestor. Ex. - forelimbs of a frog, lizard, bird and man.		$\frac{1}{2}$ $\frac{1}{2}$	
	Analogous organs – show adoption of organs for common use. Ex. – wings of butterfly and wings of bat.		$\frac{1}{2}$ $\frac{1}{2}$	
	Fossils – provide the missing links between two species. Ex. – Archeopteryx / fossils of some dinosaurs with feathers.		$\frac{1}{2}$ $\frac{1}{2}$	3
Q16.	The Sun appears reddish at Sunrise as well as at the Sunset and appears white at noon when it is overhead. At Sunrise/Sunset the blue light of shorter wave lengths, get scattered away while passing through the thicker layer of air/ large distance through air, and red light of longer wave length reach our eye. At noon, the Sun is overhead and the light of the Sun travels relatively shorter distance to our eyes and little of blue and violet colors are scattered, and white light reaches our eye.		1 1 1	 3
Q17.	<ul style="list-style-type: none"> • $h = +5 \text{ cm}$ $f = -10 \text{ cm}$ $u = -20 \text{ cm}$ $v = ?$ $h' = ?$ $\frac{1}{f} = \frac{1}{v} - \frac{1}{u}$		$\frac{1}{2}$	


	$\therefore \frac{1}{v} = \frac{1}{f} + \frac{1}{u} = \frac{1}{(-10)} + \frac{1}{(-20)} = \frac{-2-1}{20} = \frac{-3}{20}$ $\therefore v = -\frac{20}{3} \text{ cm}$ <p>Nature – Virtual, erect</p> $h' = \frac{v}{u} \times h = \left(\frac{-20}{3} \times \frac{1}{-20} \times 5 \right) \text{ cm} = +\frac{5}{3} \text{ cm}$	1 ½	
Q18.	<p>Biodegradable substances – can be broken down into simpler substances by nature / decomposers/ bacteria/ saprophytes/ saprobionts. Ex. – Human Excreta/ Vegetable peels, etc. (any one)</p> <p>Non-biodegradable substances – can't be broken down into simpler substances by nature / decomposers. Ex. – Plastic/ glass (or any other) (any one)</p> <p>Habits:</p> <ul style="list-style-type: none"> - Use of separate dustbins for biodegradable and non biodegradable waste, - Reuse of things such as poly-bags, etc., - Recycle of waste - Use of cotton /jute bags for carrying vegetables etc. <p style="text-align: right;">(any two)</p>	½ ½ ½ ½	3
Q19.	<ul style="list-style-type: none"> • Ciliary muscles modify the curvature of the eye lens to enable the eye to focus objects at varying distances/ help in adjusting the focal length of the eye lens • Presbyopia • Bifocal lens <p>(a) Defect – Myopia/ Nearsightedness Corrective lens – Concave/ Diverging lens</p> <p>(b) Values – Concerned, Caring etc. (one value of teacher, one value of Salman)</p> <p>(c) By thanking the teacher and Salman</p>	1 ½ ½ ½ ½ ½, ½ 1	3 5
Q20.	<ul style="list-style-type: none"> • Power of lens:- Ability of a lens to converge or diverge the light rays falling on it/ The degree of convergence or divergence of light rays achieved by a lens/ Reciprocal of focal length of the lens. • 1 dioptre – It is the power of a lens whose focal length is 1 metre. • $f_A = +10 \text{ cm} = 0.1 \text{ m}$ Converging/ Convex lens $P_A = \frac{1}{f_A} = \frac{1}{+0.1 \text{ m}} = +10 \text{ D}$ $f_B = -10 \text{ cm} = -0.1 \text{ m}$ <p>Diverging/ Concave lens</p> $P_B = \frac{1}{f_B} = \frac{1}{-0.1 \text{ m}} = -10 \text{ D}$ <ul style="list-style-type: none"> • In this case the object will be between the optical centre and principal 	1 ½ ½ ½ ½ ½	

focus of the lens. Hence the convex lens, i.e., lens A will form virtual and magnified image of the object.



Q21.

• Yes



(Note: image must be between F_2 and $2F_2$)

• $h = 4 \text{ cm}$ $f = +20 \text{ cm}$ $u = -15 \text{ cm}$ $v = ?$ $h' = ?$

$$\frac{1}{f} = \frac{1}{v} - \frac{1}{u}$$

$$\therefore \frac{1}{v} = \frac{1}{f} + \frac{1}{u} = \frac{1}{(+20)} + \frac{1}{(-15)} = \frac{3-4}{60} = \frac{-1}{60}$$

$$\therefore v = -60 \text{ cm}$$

Nature – Virtual, erect

$$h' = \frac{v}{u} \times h = \frac{-60 \text{ cm}}{-15 \text{ cm}} \times (+4 \text{ cm}) = +16 \text{ cm}$$

Note: Problem can be solved through ray diagram also.

Q22.

a) When Mendel cross pollinated pure tall pea plants with pure dwarf pea plants, only tall plants were obtained in F1 generation. On self pollinating the F1 progeny, both tall and dwarf plants appeared in F2 generation in the ratio 3:1

Appearance of tall character in both the F1 and F2 shows that it is a dominant character. The absence of dwarf character in F1 generation and its reappearance in F2 shows dwarfness is the recessive character.

b) When Mendel conducted a dihybrid cross having two sets of characters, he obtained only one set of parental characters in F1 generation whereas in F2 generation he obtained both the set of parental characters now recombined in the ratio of 9:3:3:1.

The appearance of new recombinants in the F2 generation along with parental type shows that traits are inherited independently.

Or

Flow chart with explanation.

1 5

½

1 ½

½

1
½

1 5

1


1

1

1

1

5

Q23.	a) Testis – secrete male hormone – testosterone	1	
	Functions – i) formation of sperms, ii) development of secondary sexual characters.	$\frac{1}{2} \times 2$	
	b) i) fallopian tube / oviduct.		
	ii) uterus.	$\frac{1}{2} \times 2$	
	• Placenta is a special disc like tissue embedded in the mother's uterine wall and connected to the foetus / embryo.	1	
	• Placenta provides a large surface area for glucose and oxygen/nutrients to pass from the mother's blood to the embryo/ foetus.	1	5
Q24.	• Soaps are sodium or potassium salts of long chain carboxylic acids.	$\frac{1}{2}$	
	• Detergents are ammonium or sulphonate salts.	$\frac{1}{2}$	
	• Cleansing action of soap – One part of soap molecule is ionic / hydrophilic and dissolves in water. The other part is non-ionic / carbon chain / hydrophobic part which dissolves in oil.	1	
	• Thus soap molecules arrange themselves in the form of a micelle / diagram of a micelle.	$\frac{1}{2}$	
	On rinsing with water, soap is washed off, lifting the oily dirt particles with it.	$\frac{1}{2}$	
	Soap does not form lather in hard water because of the reaction of soap with Ca and Mg ions present in hard water which forms insoluble ppt. / scum.	1	
	Problems due to the use of detergents are:		
	• Detergents are non-biodegradable.		
	• It leads to water or soil pollution.		
	• It can also cause skin problems. (any two)	$\frac{1}{2} \times 2$	5
SECTION – B			
	25) D	26) B	27) B
	28) D	29) B	30) C
	31) D	32) A	33) A
			1×9
Q34.	Fine	1	
		1	2
Q35.	• Carbon dioxide / CO ₂ .	1	
	• Lime Water turns milky when CO ₂ is passed through it. / A burning splinter gets extinguished by CO ₂ .	1	2

Q36. | • Towards the lens
• Magnification decreases

| 1 |
| 1 | 2